

Transferrin Monoclonal Antibody(10G3)

Description

Product type	Primary Antibody	
Code	BT-MCA1219	
Host	Mouse	
Isotype	IgG	
Size	20ul, 50ul, 100ul	
Immunogen	Protein	
Mol wt	N/A	
Species reactivity	Human	
Clonality	Monoclonal	
Recommended application	WB, IHC-p, IF, ELISA	
Concentration	l mg/ml	
Full name	Serotransferrin	
Synonyms	TF; Serotransferrin; Transferrin; Beta-1 metal-binding globulin; Siderophilin	

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

Background

This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum.


Recommended Dilution

IHC: 1:100-200 WB: 1:2000-5000 Not yet tested in other applications.


Images

94KD	-	
66KD		
45KD	- 1992	
35KD		
26KD		
	Human serum	

Western blot analysis of Human Serum with Transferrin Mouse Monoclonal antibody diluted at 1:2.000.



Immunohistochemical analysis of paraffin-embedded human-liver using antibody diluted at 1:50.



Immunohistochemical analysis of paraffin-embedded rat-liver using antibody diluted at 1:50.



501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com