

## Transferrin Monoclonal Antibody(5A4)

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-MCA1226
<b>Host</b>	Mouse
<b>Isotype</b>	IgG
<b>Size</b>	20ul, 50ul, 100ul
<b>Immunogen</b>	Protein
<b>Mol wt</b>	N/A
<b>Species reactivity</b>	Human
<b>Clonality</b>	Monoclonal
<b>Recommended application</b>	WB, IHC-p, IF, ELISA
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	Serotransferrin
<b>Synonyms</b>	TF; Serotransferrin; Transferrin; Beta-1 metal-binding globulin; Siderophilin

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

### Background

This gene encodes a glycoprotein with an approximate molecular weight of 76.5 kDa. It is thought to have been created as a result of an ancient gene duplication event that led to generation of homologous C and N-terminal domains each of which binds one ion of ferric iron. The function of this protein is to transport iron from the intestine, reticuloendothelial system, and liver parenchymal cells to all proliferating cells in the body. This protein may also have a physiologic role as granulocyte/pollen-binding protein (GPBP) involved in the removal of certain organic matter and allergens from serum.


### Recommended Dilution

IHC: 1:100-200

WB: 1:2000-5000

Not yet tested in other applications.

### Images



Western blot analysis of Human Serum with Transferrin Mouse Monoclonal antibody diluted at 1:2,000.

### Storage

-20°C for one year