

Vimentin Monoclonal Antibody

Description

Product type	Primary Antibody
Code	BT-MCA1286
Host	Mouse
Isotype	IgG
Size	50ul, 100ul
Immunogen	Purified recombinant fragment of Vimentin (aa2-466) expressed in E. Coli.
Mol wt	N/A
Species reactivity	Human,Monkey
Clonality	Monoclonal
Recommended application	WB, IHC-p, IF, ELISA
Concentration	1 mg/ml
Full name	Vimentin
Synonyms	VIM; Vimentin

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.


Background

This gene encodes a member of the intermediate filament family. Intermediate filamentents, along with microtubules and actin microfilaments, make up the cytoskeleton. The protein encoded by this gene is responsible for maintaining cell shape, integrity of the cytoplasm, and stabilizing cytoskeletal interactions. It is also involved in the immune response, and controls the transport of low-density lipoprotein (LDL)-derived cholesterol from a lysosome to the site of esterification. It functions as an organizer of a number of critical proteins involved in attachment, migration, and cell signaling. Mutations in this gene causes a dominant, pulverulent cataract.

Recommended Dilution

ELISA: 1:10000 IHC: 1:200 - 1:1000 WB: 1:500 - 1:2000 Not yet tested in other applications.

Images



Western Blot analysis using Vimentin Monoclonal antibody against HeLa (1) COS (2) HEK293 (3) and U20S (4) cell lysate.

Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, showing cytoplasmic localization with DAB staining using Vimentin Monoclonal antibody.

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com