

## Vimentin Monoclonal Antibody(1A7)

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-MCA1287
<b>Host</b>	Mouse
<b>Isotype</b>	IgG
<b>Size</b>	20ul, 50ul, 100ul
<b>Immunogen</b>	Recombinant Protein of Vimentin
<b>Mol wt</b>	53652
<b>Species reactivity</b>	Human,Mouse,Rat
<b>Clonality</b>	Monoclonal
<b>Recommended application</b>	WB
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	Vimentin
<b>Synonyms</b>	VIM; Vimentin

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

### Background


This gene encodes a member of the intermediate filament family. Intermediate filaments, along with microtubules and actin microfilaments, make up the cytoskeleton. The protein encoded by this gene is responsible for maintaining cell shape, integrity of the cytoplasm, and stabilizing cytoskeletal interactions. It is also involved in the immune response, and controls the transport of low-density lipoprotein (LDL)-derived cholesterol from a lysosome to the site of esterification. It functions as an organizer of a number of critical proteins involved in attachment, migration, and cell signaling. Mutations in this gene causes a dominant, pulverulent cataract.

### Recommended Dilution


WB: 1:1000-3000

Not yet tested in other applications.

### Images



Western blot analysis of 1) HeLa, 2) Mouse Brain Tissue, 3) Rat Brain tissue diluted at 1:2000.



Western blot analysis of 293T cell with Mouse Monoclonal antibody diluted at 1:1000

## Storage

-20°C for one year

501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China

Tel: 86 21 31007137 | E-mail: [save@bt-laboratory.com](mailto:save@bt-laboratory.com) | [www.bt-laboratory.com](http://www.bt-laboratory.com)