

## Phospho-ATF-2 (T69) Polyclonal Antibody

### Description

<b>Product type</b>	Primary Antibody
<b>Code</b>	BT-PHS00025
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Size</b>	20ul, 50ul, 100ul
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human ATF2 around the phosphorylation site of Thr69 or 51. AA range:36-85
<b>Mol wt</b>	52277
<b>Species reactivity</b>	Human, mouse, rat
<b>Clonality</b>	Polyclonal
<b>Recommended application</b>	WB, IHC-p, IP, ELISA
<b>Concentration</b>	1 mg/ml
<b>Full name</b>	Phospho-ATF-2 (T69) Antibody
<b>Synonyms</b>	ATF2; CREB2; CREBP1; Cyclic AMP-dependent transcription factor ATF-2; cAMP-dependent transcription factor ATF-2; Activating transcription factor 2; Cyclic AMP-responsive element-binding protein 2; CRE

**This product is for research use only, not for use in human, therapeutic or diagnostic procedure.**

### Background

ATF2 encodes a transcription factor that is a member of the leucine zipper family of DNA binding proteins. The activating transcription factor 2 has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. This protein binds to the cAMP-responsive element (CRE), an octameric palindrome. It forms a homodimer or a heterodimer with c-Jun and stimulates CRE-dependent transcription. This protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 *in vitro*; thus it may represent a class of sequence-specific factors that activate transcription by direct effects on chromatin components. The encoded protein may also be involved in cell's DNA damage response independent of its role in transcriptional regulation. Several alternatively spliced transcript variants have been found for this gene.

### Recommended Dilution

WB: 1: 500 - 1: 2000

IHC: 1: 100 - 1: 300

IP: 2 - 5 ug: mg lysate

ELISA: 1: 20000

Not yet tested in other applications.

### Images

No images.

### Storage

-20°C for one year