



Optimize Your Research

Recombinant Staphylokinase protein

Code CD00013

Storage: This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Intended Use

This product is for research use only, not for use in diagnosis procedures. It is highly recommended to read this instruction entirely before the use.

Source

Escherichia coli.

Purity

> 97% by SDS-PAGE and HPLC analyses.

Biological Activity

Fully biologically active when compared to standard. The specific activity determined by fibrinolytic activity in agarose plate is 5.0×10^4 IU/mg.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.

Sequence

SSSFDKGGKYK KGDDASYFEP TGPYLMVNVV GVDGKRNELL SPRYVEFPIK PGTTLTKEKI EYYVEWALDA TAYKEFRVVE LDPSAKIEVT
YYDKNKKKKEE TKSFPITEKG FVVPDLSEHI KNPGFNLITK VVIEKK

Endotoxin

Less than 1 EU/µg of rSAK as determined by LAL method.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

If you have any question on order please contact us via: order@bt-laboratory.com; technical assistance please contact us via: support@bt-laboratory.com More product visit www.bt-laboratory.com

5/F 2 Bldg., 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China
Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com