



Optimize Your Research

Recombinant Human Interleukin-17F protein

Code CD01549

Storage: This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Intended Use

This product is for research use only, not for use in diagnosis procedures. It is highly recommended to read this instruction entirely before the use.

Source

Escherichia coli.

Molecular Weight

Approximately 30.1 kDa, a disulfide-linked homodimer of two 134 amino acid polypeptide chains.

Purity

> 95 % by SDS-PAGE and HPLC analyses.

Biological Activity

Fully biologically active when compared to standard. The ED50 as determined by inducing IL-6 secretion of Mouse NIH/3T3 cells is less than 20 ng/ml, corresponding to a specific activity of $> 5.0 \times 10^4$ IU/mg.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.2, with trehalose.

Sequence

MRKIPKVGHT FFQKPESCPP VPGGSMKLDI GIINENQRVS MSRNIESRST SPWNYTVTWD PNRYPSEVVQ AQCRLGGIN
AQQKEDISMN SVPIQETLV VRRKHQGSV SFQLEKVLVT VGCTCVTPVI HHVQ

Endotoxin

Less than 1 EU/µg of rHuIL-17F as determined by LAL method.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 4 mM HCl to a concentration of 0.1mg/ml. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

If you have any question on order please contact us via: order@bt-laboratory.com; technical assistance please contact us via: support@bt-laboratory.com More product visit www.bt-laboratory.com