



Optimize Your Research

Recombinant Human Erythropoietin-Fc Chimera protein

Code CD01853

Storage: This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Intended Use

This product is for research use only, not for use in diagnosis procedures. It is highly recommended to read this instruction entirely before the use.

Source

CHO

Molecular Weight

Recombinant Human EPO/Fc produced in CHO is a dimeric, glycosylated, polypeptide chain consisting of two mature human EPO molecules linked to the Fc portion of human IgG1. The Fc component contains the CH2 domain, the CH3 domain and hinge region, but not the CH1 domain of IgG1. As a result of glycosylation, the recombinant protein migrates with an apparent molecular mass of 140 kDa in non-reducing SDS-PAGE.

Purity

> 98 % by SDS-PAGE and HPLC analyses.

Biological Activity

Fully biologically active when compared to standard. The Specific Activity was measured by the stimulation of reticulocyte production in normocyth-aemic mice and is no less than 5.0×10^5 IU/mg.

Physical Appearance

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation

Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.

Endotoxin

Less than 1 EU/µg of rHuEPO-Fc as determined by LAL method.

Reconstitution

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

If you have any question on order please contact us via: order@bt-laboratory.com; technical assistance please contact us via: support@bt-laboratory.com More product visit www.bt-laboratory.com

5/F 2 Bldg., 501 Changsheng S Rd, Nanhu Dist, Jiaxing, Zhejiang, China
Tel: 86 21 31007137 | E-mail: save@bt-laboratory.com | www.bt-laboratory.com